Express Mail Label No.: EV 570 956 967 US

INFORMATION
DISCLOSURE
CLAFIONC,
Conf. No.: 3730
Group: 1617

Express Mail Label No.: EV 370 936 907 03

Applicant: Alice C. MARTINO et al

Ser.No.: 09/656 364
Filed: September 6, 2000

Conf. No.: 3730
Atty. Docket No.: Pharmacia Case 6107.N CN2

NON PATENT LITERATURE DOCUMENTS

(g)	Ĩ	
xamuner nitials	M te	(Include Author, Title, Date, Pages, Etc.)
nitials	No.	
•	AA	Von. B. ASMUSSEN et al, Drug Release in vitro from Digoxin
1		Formulations with High Bioavailability,
		Arzneimittelforschung 1980;30(12):2168-72
	AB	A.F. DAVIS et al, Effect of supersaturation on membrane
		transport: 1. Hydrocortisone acetate, Int.J. Pharm.
		(1991), $76(1-2)$, $1-8$
	AC	C. DOHERTY et al, The In-vitro pH-Dissolution Dependence
		and In-vivo Bioavailability of Frusemide-PVP Solid
1		Dispersions, J. Pharm. Pharmacol. 41:73-78 (1989).
	AD	K.M. O'DRISCOLL et al, Chlorothiazide-Polyvinylpyrrolidone
Ī		(PVP) Interactions: Influence on Membrane Permeation
		(Everted Rat Intestine) and Dissolution, Drug Development
		and Industrial Pharmacy. 8(4),547-564(1982).
	AE	H. FUJIOKA et al, Biopharmaceutical Studies on Hydantoin
	1	Derivatives. III. Physico-chemical Properties, Dissolution
		Behavior, and Bioavailability of the Molecular Compound of
		1-benzenesulfonyl-5,5-diphenylhydantoin and Anti-Pyrine,
		J. Pharm. Dyn. 1982 Jul; 5(7): 475-484.
	AF	M. FUJII et al, Dissolution and Bioavailability of
	~~	Phenobarbital in Solid Dispersion with Phosphatidylcholine
		Chem. Pharm. Bull., 39(7)1886-1888(1991)
	AG	T. HIGUCHI, Physical Chemical Analysis of Percutaneous
	AG .	Absorption Process from Creams and Ointments,
		J.Soc.Cosmet.Chem., 11(1960)85-97.
	AH	W.I. HIGUCHI et al, Drug Membrane Transport Enhancement
	١٣.,	Using High Energy Drug-Povidone Coprecipitates,
	ŀ	Proc.Int.Symp. on Povodone, pp. 71-79, 1993.
	AI	S.L. RAGHAVEN et al, Effect of cellulose polymers on
	17.	supersaturation and in vitro membrane transport of
	1	hydrocortisone acetate, Int.J.Pharm. 193(2000) 231-237
	AJ	N. KONDO et al, Improved Oral Absorption of Enteric
	^0	Coprecipitates of a Poorly Soluble Drug, J. Pharm. Sci., 83
	j	566-570 (1994)
	AK	T. LOFTSSON et al, The effect of polyvinylpyrrolidone and
1	AK	hydroxypropyl methylcellulose on HPBCD complexation of
1		hydrocortisone and its permeability through hairless mouse
1		skin, Europ. J. Pharm. Sci. 2 (1994) 297-301
	N T	T. LOFTSSON et al, The effect of water-soluble polymers or
4	AL	aqueous solubility of drugs, Int. J. Pharm. 127 (1996) 293-
•	У .	296
,,,,,,,,,,,	1 CTCN	
AMINER	SIGNA	TURE DATE CONSIDERED

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

Express Mail Label No.: EV 570 956 967 US

Applicant: Alice C. MARTINO et al INFORMATION

Ser.No.: 09/656 364 Filed: September 6, 2000

DISCLOSURE CITETION Confictor: 3730

Atty. Docket No.: Pharmacia Case 6107.N CN2

1617 Group:

NON PATENT LITERATURE DOCUMENTS

nitials*	No.	(Include Author, Title, Date, Pages, Etc.)
	AM	N.A. MEGRAB et al, Oestradiol permeation through human skir
		and silastic membrane: effects of propylene glycol and
		supersaturation, J.Control.Rel. 36 (1995) 277-294
7	AN	H. SEKIKAWA et al, Dissolution Behaviors and
1		Gastrointestinal Absorption of Phenytoin in Phenytoin-
1	1	Polyvinylpyrrolidone Coprecipitate, Chem. Pharm. Bull., 26,
1	1	(1978) 3033-3039.
	AO	H. SUZUKI et al, Some Factors Influencing the Dissolution
1		of Solid Dispersions with Nicotinamide and
		Hydroxypropylmethylcellulose as Combined Carriers, Chem.
		Pharm. Bull., 46 1015-1020 (1998).
	AP	M.T. LEDWIDGE et al, Effects of surface active
1		characteristics and solid state forms on the pH solubility
j		profiles of drug-salt systems, Int.J.Pharm., 174 (1998)
		187-200
	AQ	Abu T.M. SERAJUDDIN et al, Influence of pH on Release of
- 1		Phenytoin Sodium from Slow-Release Dosage Forms,
- 1		J.Pharm.Sci. (1993) 82(3), 306-310
	AR	H. SUZUKI et al, Comparison of Nicotinamide, Ethylurea and
	1 .	Polyethylene Glycol as Carriers for Nifedipine Solid
ł		Dispersion Systems, Chem. Pharm. Bull., 45 (1997) 1688-1693
	AS	K. YAMAMOTO et al, Dissolution Behavior and Bioavailability
į		of Phenytoin from a Ground Mixture with Microcrystalline
		Cellulose, J. Pharm. Sci. 1976, 65(10): 1484-1488
	AT	A.P. SIMONELLI et al, Inhibition of Sulfathiazole Crystal
1	1	Growth by Polyvinylpyrrolidone, J. Pharm. Sci., 59 (1970)
1.		633-638
	AU	A.P. SIMONELLI et al, Dissolution Rates of High Energy
1		Sulfathiazole-Povidone Coprecipitates II: Characterization
1		of Form of Drug Controlling Its Dissolution Rate via
1		Solubility Studies, J. Pharm. Sci., (1976) 65(3), 355-361
EXAMINER	AV	R. WALD et al, Non-Crystallinity, Supersaturation and
	1	Relative Bioavailability: Experiences with a Non-Peptidic
		HIV-Protease Inhibitor, 10th Annual Meeting of American
	1	Assoc. of Pharm. Sci., 1995, (24 pages)

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

Form PTO-1449 (Modified)

125.05/05

- - ---- .

Page